Journal of Neuroscience Methods, 42 (1992) 237–238
Elsevier Science Publishers B.V.

Author Index (Vol. 42)

Akaoka, H., Saunier, C.-F., Chergui, K., Charléty, P., Buda, M. and Chouvet, G.
Combining in vivo volume-controlled pressure microejection with extracellular unit recording, (42) 119
Allegrini, P.R., see Sauer, D., (42) 69
Allen, T.J.A. and Knight, D.E.
The use of intracellular dialysis to study signal transduction coupling in the squid giant axon, (42) 169
Amacker, H., see Sauer, D., (42) 69

Bijlani, V., see Hayaran, A., (42) 65
Brink, P.R., see Ramanan, S.V., (42) 91
Buda, M., see Akaoka, H., (42) 119

Charléty, P., see Akaoka, H., (42) 119
Chen, Y., see Gu, Y., (42) 1
Chergui, K., see Akaoka, H., (42) 119
Chouvet, G., see Akaoka, H., (42) 119
Coimbra, A., see Merighi, A., (42) 105
Crespi, F. and Möbius, C.
In vivo selective monitoring of basal levels of cerebral dopamine using voltammetry with Nafion modified (NACRO) carbon fibre micro-electrodes, (42) 149
Cruz, F., see Merighi, A., (42) 105
Czeiger, D., see White, E.L., (42) 27

Davis, L.A., see Prochazka, A., (42) 175
De Beer, F.C., Thomeer, R.T.W.M., Marani, E. and Voogd, J.
Open fetal spinal cord surgery in rats with low mortality achieved by prevention of oligohydramnios, (42) 11
Duhl, D.M., Gillespie, D.D. and Sulser, F.
Ethidium bromide fluorescence of 28S ribosomal RNA can be used to normalize samples in Northern or dot blots when analyzing small drug-induced changes in specific mRNA, (42) 211

Ebner, T.J., see Yae, H., (42) 195
Ekström, P., see Holmqvist, B.I., (42) 45
Elias, S.A., see Yae, H., (42) 195

Fagg, G.E., see Sauer, D., (42) 69
Fan, S.F., see Ramanan, S.V., (42) 91
Farrand, D.A., see Fishman, P.S., (42) 229
Fishman, P.S., Farrand, D.A., Halpern, J.L. and Latham, W.C.
A simplified method for the preparation of tetanus toxin binding fragment for neurobiology, (42) 229

Gilbey, S.L., see Marple-Horvat, D.E., (42) 163
Gillespie, D.D., see Duhl, D.M., (42) 211
Gu, Y., Chen, Y. and Ye, L.
Electron microscopical demonstration of horseradish peroxidase by use of tetramethylbenzidine as chromogen and sodium tungstate as stabilizer (TMB-ST method): a tracing method with high sensitivity and well preserved ultrastructural tissue, (42) 1

Halpern, J.L., see Fishman, P.S., (42) 229
Hanin, I., see Zhang, X., (42) 139
Hayaran, A. and Bijlani, V.
Polyacrylamide as an infiltrating and embedding medium for vibratome sectioning of human fetal cerebellum containing DiI-filled axons, (42) 65
Hedwig, B. and Knepper, M.
Separation of synaptic and spike activity in intracellular recordings for selective analysis, (42) 83
Hirono, J., Sato, T., Tonoike, M. and Takebayashi, M.
Simultaneous recording of $[Ca^{2+}]_i$ increases in isolated olfactory receptor neurons retaining their original spatial relationship in intact tissue, (42) 185
Holmqvist, B.I., Östholm, T. and Ekström, P.
DiI tracing in combination with immunocytochemistry for analysis of connectivities and chemoarchitectonics of specific neural systems in a teleost, the Atlantic salmon, (42) 45

Jiang, W., see Pearson, K.G., (42) 75
Jirikowski, G.F.
A non-surgical technique for accurate intracerebral injections in rat, (42) 115

Kamiya, H., Sawada, S. and Yamamoto, C.
An empirical test for the reliability of quantal analysis based on Pascal statistics, (42) 19
Knepper, M., see Hedwig, B., (42) 83
Knight, D.E., see Allen, T.J.A., (42) 169
Krauth, J.
Parametric analysis of sojourn times in conditioned place preference experiments, (42) 219

Latham, W.C., see Fishman, P.S., (42) 229
Lipinski, H.-G.
Brain tissue slice thickness monitored by ion-profile measurement, (42) 37

Marani, E., see De Beer, F.C., (42) 11
Marple-Horvat, D.E. and Gilbey, S.L.
A method for automatic identification of periods of muscular activity from EMG recordings, (42) 163
Massieu, L., see Sauer, D., (42) 69
Merighi, A., Cruz, F. and Coimbra, A.
Immunocytochemical staining of neuropeptides in terminal arborization of primary afferent fibers anterogradely labeled and identified at light and electron microscopic levels, (42) 105
Möbius, C., see Crespi, F., (42) 149

Östholm, T., see Holmqvist, B.I., (42) 45

Pearson, K.G., Jiang, W. and Ramirez, J.M.
The use of naloxone to facilitate the generation of the locomotor rhythm in spinal cats, (42) 75
Prochazka, A. and Davis, L.A.
Clinical experience with reinforced, anchored intramuscular electrodes for functional neuromuscular stimulation, (42) 175

Ramanan, S.V., Fan, S.F. and Brink, P.R.
Model invariant method for extracting single-channel mean open and closed times from heterogeneous multichannel records, (42) 91
Ramirez, J.M., see Pearson, K.G., (42) 75

Sato, T., see Hirono, J., (42) 185
Sauer, D., Allegrini, P.R., Thedinga, K.H., Massieu, L., Amacker, H. and Fagg, G.E.
Evaluation of quinolinic acid induced excitotoxic neurodegeneration in rat striatum by quantitative magnetic resonance imaging in vivo, (42) 69
Saunier, C.-F., see Akaoka, H., (42) 119
Sawada, S., see Kamiya, H., (42) 19
Stein, R.B., see Wheatley, M., (42) 129
Sulser, F., see Duhl, D.M., (42) 211

Takebayashi, M., see Hirono, J., (42) 185
Thedinga, K.H., see Sauer, D., (42) 69
Thomeer, R.T.W.M., see De Beer, F.C., (42) 11
Tonoike, M., see Hirono, J., (42) 185

Voogd, J., see De Beer, F.C., (42) 11

Weinfeld, E., see White, E.L., (42) 27
Wheatley, M. and Stein, R.B.
An in vitro preparation of the mudpuppy for simultaneous intracellular and electromyographic recording during locomotion, (42) 129
White, E.L., Czeiger, D. and Weinfeld, E.
A simplified approach to retrograde/anterograde axonal labeling using combined injections of horseradish peroxidase and ibotenic acid, (42) 27
Wulfert, E., see Zhang, X., (42) 139

Yae, H., Elias, S.A. and Ebner, T.J.
Deblurring of 3-dimensional patterns of evoked rat cerebellar cortical activity: a study using voltage-sensitive dyes and optical sectioning, (42) 195
Yamamoto, C., see Kamiya, H., (42) 19
Ye, L., see Gu, Y., (42) 1

Zhang, X., Wulfert, E. and Hanin, I.
Development of a sensitive and inexpensive micropush–pull technique for the continuous analysis of brain neurotransmitters and metabolites in vivo, (42) 139

Journal of Neuroscience Methods, 42 (1992) 239–240
Elsevier Science Publishers B.V.

Key Word Index (Vol. 42)

Adenylate cyclase, (42) 169
cAMP, (42) 169
Anterograde tracing, (42) 105
Averaging, (42) 83
Axonal transport, (42) 229

Bicuculline, (42) 119

Calcium, (42) 169
Carbocyanine dyes, (42) 45
Catecholamines, (42) 139
Cerebellum, (42) 195
CGP 40116, (42) 69
Channels, (42) 91
Choline acetyltransferase, (42) 69
Chronic perfusion, (42) 139
Clonidine, (42) 75
Closed time, (42) 91
Conditioned place preference, (42) 219
Cord transection, (42) 11
Cortex, (42) 27
Cytosolic-free calcium, (42) 185

Diaminobenzidine, (42) 1
DiI, (42) 65
Dopamine, (42) 149
Dorsal horn, (42) 105
Double labeling, (42) 45
Dye recovery, (42) 139

Electrode fabrication, (42) 119
Electron microscopy, (42) 1; (42) 105
Embedding, (42) 65
EMG – automated analysis, (42) 163
Exponential model, (42) 219
Extracellular diffusion, (42) 37

Fetal rats, (42) 11
Fragment C, (42) 229
Fura-2, (42) 185

GABA, (42) 119
Glutamic acid decarboxylase, (42) 69
GTP-binding proteins, (42) 169
Guide cannula, (42) 139

[^{3}H]Choline recovery, (42) 139
Heterogeneity, (42) 91
Hippocampus, (42) 139
Horseradish peroxidase, (42) 1; (42) 27; (42) 105
Human fetus, (42) 65
Hypothalamus, (42) 115

Ibotenic acid, (42) 27
Image restoration, (42) 195
Immunocytochemistry, (42) 105
Immunogold, (42) 105
Implanted functional electrical stimulation electrodes, (42) 175
Infiltration, (42) 65
Injection aid, (42) 115
Internal standards, (42) 211
Interneuron, (42) 129
Intracellular recordings, (42) 83
Intracerebral injection, (42) 115
Invariance, (42) 91
In vivo voltammetry, (42) 149
Ion profile, (42) 37
Ion-selective microelectrode, (42) 37
Isolation, (42) 185

Lateral ventricle, (42) 139
Locomotion, (42) 75; (42) 129
Locus coeruleus, (42) 119

Maximum likelihood estimator, (42) 19
Mean times, (42) 91
Microperfusion, (42) 139
Monte Carlo simulation, (42) 19
Motor control, (42) 129
Multiple labeling, (42) 105
Mycoplasma infection, (42) 11

NA-CRO micro-biosensor, (42) 149
Naloxone, (42) 75
Neurodegeneration, (42) 69
Neuronal labeling, (42) 229
NMDA receptor, (42) 69
N-methyl D-aspartate, (42) 129
Non-stationarity, (42) 19
Normalization, (42) 211
Northern blots, (42) 211

Olfactory receptor, (42) 185
Olfactory system, (42) 45
Oligohydramnios, (42) 11

Open surgery, (42) 11
Open time, (42) 91
Optical imaging, (42) 195
Optical recording, (42) 185
Optical sectioning, (42) 195
Optic tectum, (42) 45

Pascal distribution, (42) 19
Pattern generation, (42) 75
Pattern generator, (42) 129
Pineal organ, (42) 45
Pituitary, (42) 45
Polyacrylamide, (42) 65
Post-embedding, (42) 105
Postsynaptic potentials, (42) 83
Pressure microejection, (42) 119
Projection, (42) 27
Purkinje cells, (42) 195

Quantal analysis, (42) 19
Quantitation, (42) 211

Reciprocal, (42) 27
Retina, (42) 45

Salmo salar L, (42) 45
Serotonin (5-HT), (42) 169
Signal transduction coupling, (42) 169
Single-unit recording, (42) 119
Slice swelling, (42) 37
Slice thickness, (42) 37
Sodium tungstate, (42) 1
Sojourn times, (42) 219
Spike filter, (42) 83
Spinal cord, (42) 129
Squid axon, (42) 169
28S RNA, (42) 211
Striatum, (42) 149
Substantia nigra, (42) 119
Synapse, (42) 19

Tetramethylbenzidine, (42) 1
Thalamus, (42) 27
Tissue printing, (42) 185
Transmitter release, (42) 19
Transneuronal transport, (42) 229

Voltage-sensitive dye, (42) 195

Walking, (42) 75